Resource for Unit 1 Lesson 2- The Epidermis

Lesson 2 The Epidermis.

This is the topmost layer of the skin. Within the epidermis there are 5 layers.

Let's start with the top most outer layer of the epidermis- called the stratum corneum

Stratum Corneum

The stratum corneum serves as the outermost layer of the skin and plays a pivotal role in maintaining the body's integrity through its distinct structure and functions. This sophisticated structure, with its intricate composition and dynamic properties, is of paramount importance for both its protective barrier function and its involvement in various physiological processes.

At a cellular level, the stratum corneum is primarily composed of **corneocytes**, flattened **keratinocytes** that have undergone terminal differentiation. These corneocytes are held together by an intercellular matrix comprised of lipids, proteins, and natural moisturizing factors. This intricate matrix contributes significantly to the barrier properties of the stratum corneum.

One of the defining features of the stratum corneum is its "brick and mortar" arrangement, wherein the corneocytes act as the "bricks" and the intercellular lipids serve as the "mortar." This arrangement creates a highly impermeable barrier, effectively preventing excessive water loss, environmental pollutants, and pathogens from entering the body while simultaneously minimizing the escape of essential water and electrolytes.

The stratum corneum plays a pivotal role in maintaining skin hydration. Natural moisturizing factors (NMFs) within the stratum corneum, including amino acids, urea, and lactic acid, contribute to the skin's ability to attract and retain water, thereby maintaining optimal hydration levels. This function is vital not only for skin health but also for overall homeostasis.

Amino Acids: Amino acids are the building blocks of proteins. NMFs in the skin often include amino acids such as arginine, serine, and histidine. These amino acids help bind water molecules, preventing excessive water loss from the skin.

Hygroscopic Compounds: Certain substances in the NMFs are hygroscopic, meaning they have a strong affinity for water. These include urea and lactic acid. They attract moisture from the environment and retain it in the skin, contributing to its hydration.

Inorganic Ions: NMFs also contain inorganic ions like sodium, potassium, and calcium. These ions help regulate the osmotic balance of the skin, which is critical for maintaining proper hydration levels.

Natural Peptides: Peptides derived from enzymatic breakdown of larger proteins can also be found in NMFs. These peptides can have moisturizing and soothing effects on the skin.

Urocanic Acid: Urocanic acid is a breakdown product of histidine and is involved in regulating the skin's pH. A balanced pH is important for maintaining skin health and hydration.

Pyrrolidone Carboxylic Acid (PCA): PCA is a derivative of glutamic acid and has a strong ability to bind water. It helps to maintain skin moisture and acts as a humectant.

NMFs are produced by the skin's natural metabolic processes, and their levels can be influenced by various factors, including genetics, age, and environmental conditions. When the NMF content is disrupted or depleted, it can lead to dry, dehydrated skin, making it a significant consideration in skincare and aesthetic practices.

In skincare routines and aesthetic treatments, products containing NMF-like substances or ingredients that support the skin's natural ability to maintain moisture are often used. These can include moisturizers, serums, and treatments designed to enhance the skin's hydration and barrier function.

Understanding NMFs and their role in skin health is essential for professionals in the field of aesthetics, as it informs the selection of appropriate skincare products and treatments to address specific skin concerns related to hydration and moisture retention

The stratum corneum exhibits dynamic properties that respond to various external stimuli. For instance, its permeability barrier can be modulated by factors such as humidity, temperature, and pH, which influence the organization of the lipid matrix. This adaptability is crucial for the skin's ability to adjust to changing environmental conditions.

Individual Components of the Stratum Corneum

Corneocytes: At the heart of the stratum corneum are corneocytes, which are flattened, enucleated keratinocytes that have undergone terminal differentiation. These specialized cells are held together in a cohesive manner, forming the sturdy "brick" component of the barrier. Corneocytes are rich in keratin proteins, which provide mechanical strength to the skin.

Intercellular Lipids: The "mortar" that holds the corneocytes together is composed of a complex mixture of lipids, including ceramides, cholesterol, and free fatty acids. These lipids form bilayers within the extracellular spaces, creating a barrier that effectively prevents the entry of water, pathogens, and foreign substances. The composition and organization of these lipids are crucial for maintaining the barrier's integrity and function.

Natural Moisturizing Factors (NMFs): NMFs are a group of small molecules, primarily amino acids, urea, and lactic acid, that contribute to the skin's ability to retain water. They play a vital role in maintaining skin hydration by attracting and binding water molecules, thus ensuring the suppleness and flexibility of the skin. NMFs also contribute to the regulation of the skin's pH.

Proteases and Protease Inhibitors: Enzymes known as proteases are present in the stratum corneum, and their primary role is to break down the bonds between corneocytes. This controlled degradation of corneocytes facilitates the shedding of dead skin cells from the surface, allowing for the continuous renewal of the skin. Protease inhibitors help regulate this process to prevent excessive breakdown.

Desmosomes: Desmosomes are specialized cell junctions that connect adjacent corneocytes, adding structural stability to the stratum corneum. These adhesive structures contribute to the cohesive nature of the skin's surface and prevent the corneocytes from separating too easily.

Langerhans Cells: While not as prominent as other components, Langerhans cells are immune cells found within the stratum corneum. These dendritic cells play a role in detecting and responding to potential pathogens, thereby contributing to the skin's immune defense.

Cornified Envelope: This is a protein structure that forms just beneath the cell membrane of corneocytes. It helps in maintaining the cell's shape and protecting it from external stresses. The cornified envelope is formed through the cross-linking of various proteins, including involucrin, loricrin, and filaggrin.

Barrier Lipids: Beyond the intercellular lipids, specific barrier lipids, such as ceramides, are crucial for maintaining the stratum corneum's barrier function. These lipids contribute to the cohesion of the lipid matrix and play a role in regulating transepidermal water loss (TEWL).

Epidermal Lipid Processing: Enzymes within the stratum corneum are responsible for converting precursor molecules into the lipid constituents needed for the barrier. This processing is essential for maintaining the balance of lipids and ensuring the proper function of the barrier.

In summary, the stratum corneum is a composite structure comprised of corneocytes, intercellular lipids, natural moisturizing factors, proteases, desmosomes, Langerhans cells, the cornified envelope, barrier lipids, and enzymes involved in lipid processing. Each of these components contributes uniquely to the barrier's protective function, the skin's hydration balance, immune response, and overall skin health.

Stratum Lucidim

The stratum lucidum, often referred to as the "clear layer," is a distinctive and less commonly present layer. It appears in certain areas of the skin, such as the palms of the hands and the soles of the feet. It is situated between the stratum granulosum and the stratum corneum and serves a specific purpose in enhancing the skin's resilience and protection in those areas.

Characterized by its translucent appearance under a microscope, the stratum lucidum is composed of flattened, densely packed keratinocytes that lack nuclei and other organelles. This layer is notable for its high keratin content, which contributes to its transparent appearance and the protective properties of the skin in regions subject to constant mechanical stress and friction.

The primary function of the stratum lucidum is to provide an additional layer of mechanical strength to the skin. In areas that experience repeated pressure and friction, such as the palms and soles, this layer acts as a barrier against wear and tear. The densely packed keratinocytes in the stratum lucidum are tightly interconnected through **desmosomes**, specialized cell junctions that enhance the layer's integrity and resilience.

Desmosomes are specialized cell junctions found in the skin and other tissues. They play a crucial role in maintaining the structural integrity and stability of the epidermis, which is the outermost layer of the skin. In your role as an aesthetic professional, understanding desmosomes is valuable because they contribute to the overall health and appearance of the skin. Here's a detailed explanation of desmosomes:

Desmosomes are a type of cell-to-cell junction that functions to firmly anchor adjacent cells together, providing mechanical strength and preventing them from easily separating. In the context of the skin, desmosomes are particularly important in the epidermal layers, where cells are constantly exposed to mechanical stress, such as stretching and friction.

Key features and functions of desmosomes in the skin include:

Cell Adhesion: Desmosomes are composed of specialized proteins known as desmogleins and desmocollins, which extend from the cell membrane into the cytoplasm. These proteins interact with similar proteins on adjacent cells, forming strong connections that bind the cells together. This adhesion is essential for maintaining tissue integrity.

Intermediate Filaments: Inside the cell, desmosomes connect to intermediate filaments made of a protein called keratin. These intermediate filaments provide structural support and help distribute mechanical stress across the cells, reducing the risk of cell damage or separation.

Tissue Strength: Desmosomes contribute to the overall strength of the epidermis, allowing it to withstand various stresses, including bending, stretching, and compression. Without desmosomes, the skin would be more fragile and susceptible to damage.

Clinical Implications: Disruptions in desmosome function can lead to skin conditions such as pemphigus and pemphigoid, which are autoimmune diseases characterized by blistering and skin fragility due to the loss of cell adhesion.

In the field of aesthetics, understanding the role of desmosomes can be relevant when considering treatments and procedures that affect the skin's integrity, such as chemical peels, microdermabrasion, or laser therapies. These procedures may temporarily disrupt desmosomal connections as part of the skin's natural healing and remodeling process. Therefore, it's important to take into account the recovery and maintenance of desmosome function when assessing post-treatment outcomes and skincare regimens.

It's important to note that the presence and thickness of the stratum lucidum can vary based on factors such as genetics and the location on the body. In areas with thicker layers of the stratum lucidum, the skin appears tougher and less permeable.

In summary, the stratum lucidum is a specialized layer within the epidermis that is found in specific regions of the body, such as the palms and soles. Composed of densely packed, flattened keratinocytes, its translucent appearance and high keratin content contribute to the skin's mechanical strength and protection against friction and external stressors. While not present in all skin regions, the stratum lucidum plays a crucial role in enhancing the skin's resilience and barrier function.

Stratum Granulosum

The stratum granulosum marks a transition between actively dividing cells and the outermost protective layer of the skin, the stratum corneum.

The stratum granulosum is characterized by the presence of **granules** within its keratinocytes, the main cells of the epidermis. These granules, known as **keratohyalin granules**, contain important proteins that contribute to the formation of the skin's protective barrier. As keratinocytes migrate

upwards from the basal layer, they undergo a transformation within the stratum granulosum that prepares them for their role in the stratum corneum.

During this process, keratinocytes accumulate keratohyalin granules, which contain proteins like **profilaggrin.** Profilaggrin is a crucial protein that plays a central role in aggregating keratin filaments, essential components for the mechanical strength of the skin. As these granules accumulate, they interact with the keratin filaments, promoting their bundling and alignment, a process that sets the stage for the formation of the cornified envelope—an essential structural component of the stratum corneum.

Simultaneously, as keratinocytes move upwards through the stratum granulosum, a process known as keratinization occurs. This involves the breakdown of organelles and the restructuring of cellular components to provide energy for the maturation process. As keratinocytes lose their nuclei and cellular organelles, they become flattened and densely packed with the keratin filaments, creating a tougher, more protective layer.

The stratum granulosum also plays a role in lipid secretion, contributing to the formation of the skin's lipid barrier. **Lamellar bodies**, specialized organelles within keratinocytes, produce lipids that are released into the extracellular space. These lipids contribute to the intercellular lipid matrix that acts as a waterproof seal, preventing the entry of harmful substances and minimizing water loss.

Individual Components in the Stratum Granulosum

Keratohyalin Granules: These specialized granules are a hallmark of the stratum granulosum. They contain important proteins, such as profilaggrin, which play a central role in the formation of the cornified envelope—a critical structural component of the stratum corneum. Profilaggrin undergoes enzymatic cleavage to produce filaggrin, a protein that promotes the bundling and alignment of keratin filaments within keratinocytes.

Filaggrin: As keratinocytes transition from the stratum granulosum to the stratum corneum, filaggrin-rich granules are broken down into free amino acids, urocanic acid, and pyrrolidone carboxylic acid. These breakdown products contribute to the skin's natural moisturizing factors (NMFs), enhancing its ability to retain water and maintain proper hydration.

Lamellar Bodies: These membrane-bound organelles are present within keratinocytes of the stratum granulosum. Lamellar bodies are responsible for the synthesis and secretion of lipids, such as ceramides, cholesterol, and free fatty acids. These lipids are crucial for forming the intercellular lipid matrix in the stratum corneum, which acts as a barrier against water loss and protects the skin from external stressors.

Lipid Bilayers and Lamellar Structures: Within the stratum granulosum, lipid bilayers and lamellar structures are formed through the secretion of lipids from lamellar bodies. These structures contribute to the cohesive lipid matrix in the stratum corneum, enhancing its barrier function and minimizing transepidermal water loss.

Cellular Organelle Degradation: As keratinocytes move from the basal layer to the stratum granulosum, their cellular organelles, including the nucleus and most cytoplasmic organelles, degrade and are eventually lost. This process is essential for the formation of the corneocytes that make up the stratum corneum, as it contributes to the mechanical strength and water-resistant properties of the outermost skin layer.

Keratinization and Cornification: The process of keratinization involves the transformation of keratinocytes into corneocytes. This process includes the accumulation of keratohyalin granules, the breakdown of cellular components, and the alignment of keratin filaments. Cornification is the final stage, in which the corneocytes lose their nuclei and cellular organelles, becoming densely packed with keratin filaments and contributing to the strength and resilience of the stratum corneum.

Regulation of Epidermal Barrier Formation: The stratum granulosum also contributes to the regulation of barrier formation through the expression of various enzymes and proteins. These components play a role in the processing of lipids, the synthesis of filaggrin, and the formation of the cornified envelope.

In summary, the stratum granulosum is a crucial layer in the epidermis that orchestrates the transformation of keratinocytes into mature corneocytes, setting the stage for the formation of the skin's protective barrier.

Stratum Spinosum or Spiny Layer.

The stratum spinosum, often referred to as the "spiny layer," that lies above the basal layer and below the stratum granulosum. It plays a pivotal role in the epidermal barrier formation, cell communication, and defence against environmental stressors.

One of the defining features of the stratum spinosum is the presence of **desmosomes,** specialized cell junctions that appear under the microscope as small, spiny projections connecting adjacent keratinocytes. These desmosomes provide mechanical strength and adhesion, holding the keratinocytes together in a cohesive manner. This feature is particularly significant for the skin's resilience and ability to withstand physical stress.

Within the stratum spinosum, keratinocytes actively synthesize keratin proteins—a key component of the skin's structure. As these cells move upward from the basal layer, they begin to differentiate and undergo changes in their cytoskeletal architecture. This differentiation process contributes to the formation of the skin's protective barrier by promoting the synthesis and accumulation of lipids and proteins that are essential for the stratum corneum's function.

The stratum spinosum also serves as a platform for cell communication and immune responses. **Langerhans cells,** specialized immune cells, are found within this layer. These cells play a crucial role in detecting and responding to potential pathogens and antigens, contributing to the skin's immune defence. Additionally, the stratum spinosum facilitates the exchange of signals between keratinocytes, allowing for coordinated responses to various environmental challenges.

So, the stratum spinosum is involved in the regulation of cell proliferation and differentiation. As keratinocytes move from the basal layer to the stratum spinosum, they continue to divide and differentiate, preparing for their roles in the outer layers of the epidermis. This controlled proliferation and differentiation ensure the continuous renewal of the epidermis and the maintenance of skin integrity.

Individual Components in the Stratum Spinosum

Desmosomes: Desmosomes are specialized cell junctions that are abundant in the stratum spinosum. These structures play a critical role in providing mechanical strength and cohesion to the

skin. Desmosomes are formed by transmembrane proteins called desmogleins and desmocollins, which interact with similar proteins on adjacent cells, effectively "stitching" the cells together. This adhesive function ensures that the layers of the epidermis remain tightly connected, even under mechanical stress.

Keratin Synthesis: Keratinocytes within the stratum spinosum are actively engaged in synthesizing keratin proteins. These intermediate filaments provide structural integrity to the skin and are responsible for its strength and resilience. As keratinocytes differentiate and move towards the surface, the synthesis of keratin proteins increases, contributing to the formation of the stratum corneum's cornified envelope.

Langerhans Cells: Langerhans cells are dendritic immune cells present within the stratum spinosum. They play a vital role in the skin's immune defence by detecting and capturing antigens or pathogens that come into contact with the skin. Langerhans cells then migrate to nearby lymph nodes to present these antigens to other immune cells, initiating an immune response when necessary.

Cell Communication and Signalling: The stratum spinosum serves as a communication hub between keratinocytes and other skin cells. Gap junctions, which are specialized channels between adjacent cells, allow for the exchange of ions and small molecules, facilitating coordinated responses to external stimuli. This communication is vital for maintaining tissue homeostasis and responding to environmental changes.

Proliferation and Differentiation: Keratinocytes in the stratum spinosum continue to divide and differentiate. This controlled cell proliferation ensures a constant supply of new cells for the upper layers of the epidermis. As keratinocytes move towards the surface, they undergo changes in their organelles and cytoskeletal structure, preparing them for their roles in the outer layers.

Involvement in Barrier Formation: While the stratum granulosum is primarily associated with barrier formation, the stratum spinosum contributes to this process as well. The synthesis of lipids and proteins in the stratum spinosum contributes to the formation of the skin's natural barrier, setting the stage for further maturation in the stratum granulosum and stratum corneum.

Cytoskeletal Changes: As keratinocytes differentiate within the stratum spinosum, they undergo changes in their cytoskeletal structure. This process involves the reorganization of cytoskeletal components, including intermediate filaments, microtubules, and microfilaments. These changes are essential for the cells' ability to adapt to different mechanical and physiological demands.

Langerhans Cells: are a specialized type of immune cell found predominantly in the skin's epidermis. These cells play a pivotal role in the body's immune response by serving as key antigen-presenting cells and contributing to the defence against pathogens and foreign substances.

Structurally, Langerhans cells are classified as dendritic cells due to their branched, tree-like appearance. They possess long, membrane-bound extensions called dendrites that extend among the epidermal keratinocytes, forming a network that enables them to survey their surroundings. Langerhans cells are characterized by the presence of Birbeck granules within their cytoplasm, which have a tennis racket-like shape and are considered a unique marker of these cells.

The primary function of Langerhans cells lies in their role as antigen-presenting cells (APCs). When the skin is exposed to various antigens, such as pathogens or allergens, Langerhans cells capture these antigens using their dendrites. Once an antigen is captured, Langerhans cells process it

internally and break it down into smaller fragments. These fragments are then presented on the surface of the Langerhans cell using specialized molecules called major histocompatibility complexes (MHC) class II molecules.

This process of antigen presentation is crucial for initiating an immune response. When a T lymphocyte (a type of white blood cell) encounters an antigen presented by a Langerhans cell, it triggers an immune reaction. This immune response can involve the activation of various immune cells, including T cells and B cells, leading to the elimination of pathogens and the generation of immune memory.

Langerhans cells also have a unique property in that they are capable of tolerogenic responses. In certain situations, Langerhans cells can induce a state of immune tolerance, preventing an excessive immune response to harmless substances. This function is important for maintaining immune balance and preventing the development of autoimmune reactions.

In summary, the stratum spinosum is a complex and dynamic layer within the epidermis that contributes to the skin's structure, immune defence, cell communication, and barrier formation. Its key components, including desmosomes, keratin synthesis, Langerhans cells, cell communication, and controlled proliferation, collectively ensure the skin's strength, resilience, and adaptability.

Basal Layer

The basal layer, also known as the stratum basale, constitutes the innermost layer of the epidermis and rests on top of the basement membrane that separates the epidermis from the underlying dermis. It is a dynamic and crucial layer responsible for continuous cell renewal, skin regeneration, and the formation of the upper epidermal layers.

Composed primarily of proliferating keratinocytes, the basal layer is characterized by its actively dividing cells. These keratinocytes are anchored to the basement membrane by **hemidesmosomes**, specialized cell junctions that provide structural stability and prevent the cells from detaching from the underlying dermis.

The basal layer plays a pivotal role in the generation of new epidermal cells through a process known as keratinocyte proliferation and differentiation. Basal keratinocytes are **stem-like** cells that give rise to daughter cells through **mitosis**. As these daughter cells are produced, they move upwards through the epidermal layers, gradually differentiating and undergoing changes in structure and composition.

One of the significant functions of the basal layer is the synthesis of keratin proteins, which are essential for the skin's mechanical strength and barrier function. As keratinocytes differentiate and move upward, they accumulate keratin proteins and undergo morphological changes that prepare them for their roles in the outer layers of the epidermis.

The basal layer houses **melanocytes**, specialized cells responsible for producing melanin—the pigment that provides skin colour and offers protection against harmful ultraviolet (UV) radiation. Melanocytes transfer melanin to neighbouring keratinocytes, contributing to the skin's UV protection and minimizing potential DNA damage caused by UV exposure.

The basal layer also contains **Merkel cells**, which are involved in sensory perception and touch sensation. These cells are found in regions of the body that are particularly sensitive to touch, such

as the fingertips and lips. Merkel cells form connections with nerve endings, aiding in the perception of tactile stimuli.

.

Individual Components in the Basal Layer

Keratinocytes: Keratinocytes are the predominant cell type within the basal layer and serve as the foundation for epidermal renewal and regeneration. Stem-like basal keratinocytes continuously divide through mitosis to produce daughter cells. These daughter cells, known as transit amplifying cells, move upward through the epidermal layers, ultimately differentiating into the specialized cells that make up the upper layers of the epidermis. Keratinocytes are responsible for producing keratin proteins, which provide the structural integrity and resilience of the skin.

Hemidesmosomes: Hemidesmosomes are specialized cell junctions that anchor basal keratinocytes to the basement membrane—the interface between the epidermis and the underlying dermis. Hemidesmosomes play a crucial role in maintaining the structural integrity of the skin and preventing the detachment of the epidermal layers from the dermis. This anchoring ensures that the epidermis remains firmly attached, even under mechanical stress.

Melanocytes: Melanocytes are specialized cells found in the basal layer that are responsible for producing melanin—the pigment responsible for skin, hair, and eye colour. Melanin also provides protection against harmful ultraviolet (UV) radiation by absorbing and dissipating UV energy. Melanocytes transfer melanin to neighbouring keratinocytes through long, branching cellular processes, contributing to the skin's UV defence and overall colouration.

Melanocytes are specialized cells located within the basal layer of the epidermis. They are primarily found in regions of the skin exposed to sunlight, such as the face, arms, and hands. Melanocytes produce the pigment melanin, which determines the colour of our skin, hair, and eyes.